

Correlations and Path Coefficient Analysis of Major Quantitative Characters in Tef [*Eragrostis tef* (Zucc.) Trotter] Breeding Lines

Habte Jifar Daba

Research Article

Department of Genetics and Plant Breeding, Ethiopian Institute of Agricultural Research, Ethiopia

*Corresponding author: Habte Jifar Daba, Department of Genetics and Plant Breeding, Ethiopian Institute of Agricultural Research, Ethiopia, Tel: +25191 780 7238; E-mail: habtejjfar@gmail.com

Received: February 20, 2019; Accepted: March 12, 2019; Published: March 19, 2019

Copyright: © 2019 Daba HJ. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Grain yield is a complex quantitative trait resulting from the interaction of various genetic and environmental factors. Knowledge of the association of characters and the direct and indirect effects of the various characters on grain yield is essential for effective selection in crop improvement. Since such studies are very limited in tef, further research is required. This study was designed to examine the interrelationship among 12 characters and their direct and indirect effect on grain yield. Thus, 28 tef genotypes were evaluated at Holetta, Debre Zeit and Alem Tena during the main cropping season of 2015 in randomized complete block design with three replications. The study results revealed significant and positive genotypic and phenotypic correlation between grain yield and all studied traits other than days to heading and to maturity, lodging index and thousand kernel weight. The estimated values of genotypic correlations were also found to be higher than their respective phenotypic correlations. In the path coefficient analysis, very high (nearly one) genotypic direct effects on grain yield per plot was exhibited by total biomass and harvest index, respectively. Total biomass, furthermore, showed the highest genotypic indirect effect on grain yield via all traits other than lodging index and thousand kernel weight. Total biomass, harvest index, second culm diameter, plant height and panicle length, in general, showed positive and significant genotypic association with grain yield. Hence, these characters could be used as target traits for the improvement of grain yield in tef.

Keywords: Direct effect; Improvement; Indirect effect; Recombinant inbred lines; Tef; Traits

Introduction

Tef [*Eragrostis tef* (Zucc.) Trotter] is the most important indigenous crops widely grown and consumed in Ethiopia. It is a crop with various agronomic, nutritional and health merits. Though tef stands first in area coverage and second to maize in total grain production, its average productivity is still very low compared to other major cereal crops grown in Ethiopia. The productivity of tef in 2016/17, for instance, is 16.5 t/ha compared to 25.3 t/ha for sorghum, 26.7 t/ha for wheat and 36.8 t/ha for maize [1]. Poor dissemination of improved varieties and production packages, and the problem of lodging are among the major factors adversely affecting the productivity of tef.

Grain yield is a complex quantitative trait resulting from the interaction of various genetic and environmental factors. Genotypic and/or phenotypic correlation coefficients are useful to measure the association among various traits of a given crop and also enable to determine components that influence a given trait either directions [2-4]. Though correlation coefficients measure the absolute value of the correlation between variables in a given body of data, it may not provide clear picture of the relative importance of direct and indirect effect of independent traits on dependent traits [4-6]. In other words, correlation measures only mutual association without considering causation while, path coefficient measures the direct influence of one variable upon another variable. Hence, path coefficient analysis permits the partitioning of correlation coefficient analysis, grain yield is usually considered to be dependent while the other traits are

considered independent. Since it permits critical estimation of specific factor that produce a given correlation, path coefficient analysis has been successfully employed in formulating an effective selection strategy for a breeding program [6,7]. Thus, generating information on the association of characters and the direct and indirect effects of each traits on grain yield are essential to aid the selection process in crop improvement. The objectives of the present study were, therefore, to estimate the phenotypic and genotypic correlations among 12 morphoagronomic traits, and to determine the direct and indirect effect of various traits on grain yield of tef.

Materials and Methods

Plant materials

Twenty-eight tef genotypes including 24 breeding lines, three improved varieties (standard checks) and one farmers' variety (local check) were evaluated at Holetta, Debre Zeit and Alem Tena in 2015 main cropping season. The studied breeding lines are entirely recombinant inbred lines (RIL). Quncho is a popular variety released for high rainfall high potential environments while, Tsedey is the name of variety released for moisture stress areas of Ethiopia. Kinde, on the other hand, is a semi-dwarf mutant line identified at the University of Bern from mutagenized tef population. Further, description of the study genotypes is given in Table 1. Regarding the study environments, Holetta and Debre Zeit are found in the central highland high rainfall area and mid altitude environment, respectively while, Alem Tena is located in the moisture deficit Rift Valley area of Ethiopia (Table 2). Citation: Habte Jifar Daba (2019) Correlations and Path Coefficient Analysis of Major Quantitative Characters in Tef [*Eragrostis tef* (Zucc.) Trotter] Breeding Lines. Adv Crop Sci Tech 7: 431.

S No.	Genotype names	Descriptio n	S No.	Genotype names	Description	8	RIL-121	Breeding line	22	RIL-159	Breeding line
1	RIL-13	Breeding line	15	RIL-171	Breeding line	9	RIL-69	Breeding line	23	RIL-85	Breeding line
2	RIL-81	Breeding line	16	RIL-91	Breeding line	10	RIL-134	Breeding line	24	RIL-137	Breeding line
3	RIL-302	Breeding line	17	RIL-115	Breeding line	11	RIL-11	Breeding line	25	Quncho	Released varie
4	RIL-232	Breeding line	18	RIL-180	Breeding line	12	RIL-133	Breeding line	26	Kinde	Released varie
5	RIL-227	Breeding line	19	RIL-103	Breeding line	13	RIL-271	Breeding line	27	Tsedey	Released varie
6	RIL-181	Breeding line	20	RIL-96	Breeding line	14	RIL-244	Breeding line	28	Local	Farmers' variet
7	RIL-110	Breeding	21	RIL-132	Breeding line	Tabl	e 1. Names o	of study genoty	mes an	d their descri	intions

Locations	Altitudo	Latitudo	Longitudo	rainfall (mm)	Temperature		Soil type
			Longitude	rannan (ninn)	Min °C	Max °C	
Alem Tena	1580	8020' N	38057'E'	500	8	29.8	Light sandy
Debre Zeit	1900	8 044' N	380 58' E	851	8.9	28.3	Pellic Vertisol
Holetta	2400	9044'N	380 30' E	1100	6	22	Nitosol

Table 2: Description of the edaphic and climatic conditions of the study locations.

The experiment was laid out in randomized complete block design with three replications. Each genotype was grown on five rows of 1.0 m length at a spacing of 0.2 m, 1 m and 1.5 m, between rows, plots and replications, respectively. All agronomic and cultural practices were applied as per the recommendation for tef production at each location. Plant height, panicle length, peduncle length, culm diameter, number of spikelet per panicle data were collected based on five randomly tagged plants before heading while days to panicle emergence, days to maturity, total biomass, grain yield, 1000-kernel weight and lodging index data were recorded on plot basis.

Statistical analysis

Before proceeding to correlation and path coefficient analysis, all quantitative traits data were subjected to analysis of variance (ANOVA) as per the suggested procedure [8]. Genotypic and phenotypic correlation analysis were then conducted using multi environment trial analysis with R for windows (META-R) version 6.0 developed by CIMMYT [9]. Genes statistical software packages version 1990.2018.39 [10] was used for path coefficient analysis while cluster analysis of the 12 studied traits was done using MINITAB software version 17.1 [11].

Results and Discussion

Correlation coefficient analysis

Estimates of genotypic and phenotypic correlations among the various traits of tef are presented in Table 3. Thus, positive and significant genotypic and phenotypic association were exhibited between grain yield and all studied traits, except days to heading and maturity, lodging index and thousand kernel weight. This suggests that genotypes heading earlier have better yield than those with longer heading date. This is contrary to most previous studies [3,5] which reported genotypes with longer heading date have higher yield than the earlier ones. On the other hand, the non-significant association detected between grain yield and days to maturity and lodging in the present study is in line with the previous report [3]. The positive association exhibited between grain yield and pant height indicates that genotypes with longer plant height have usually higher grain yield compared to those having shorter plant height. Similar findings have been reported previously [3,5]. Such observed positive correlation among the studied traits could be due to presence of common genetic elements that control the characters to the same direction.

Traits	DH	DM	РН	PL	Pdl	SCD	SPK	твм	н	LI	ткw	GY
DH	1.00	0.64b	0.35	0.45 ^c	-0.19	0.16	0.57 ^b	0.19	-0.78 ^a	0.12	-0.37 ^c	-0.17
DM	0.92 ^a	1.00	0.48 ^c	0.42 ^c	0.05	0.21	0.39 ^c	0.42 ^c	-0.40 ^c	0.16	-0.08	0.20

Page 2 of 6

PH	0.44 ^c	0.73 ^a	1.00	0.87 ^a	0.27	0.50 ^b	0.70 ^b	0.89 ^a	-0.08	0.19	-0.14	0.75 ^a
PL	0.74 ^a	0.79 ^a	1.00 ^a	1.00	0.00	0.50 ^b	0.74 ^a	0.73 ^a	-0.23	0.12	-0.06	0.55 ^b
Pdl	-0.21	0.22	0.33	0.04	1.00	0.00	-0.06	0.27	0.23	-0.01	-0.19	0.33
SCD	0.34	0.44 ^c	1.00 ^a	1.00 ^a	0.08	1.00	0.46 ^c	0.62 ^b	-0.05	-0.12	-0.04	0.52 ^b
SPK	0.91 ^a	0.72 ^b	0.81 ^a	0.95 ^a	-0.13	1.00 ^a	1.00	0.63 ^b	-0.42 ^c	0.16	-0.24	0.38 ^c
твм	0.22	0.67 ^b	1.00 ^a	1.00 ^a	0.34	1.00 ^a	0.82 ^a	1.00	0.02	0.02	-0.02	0.90 ^a
н	-1.00 ^a	-0.63 ^b	-0.14	-0.37 ^c	0.29	-0.34	-0.83 ^a	0.21	1.00	0.14	0.27	0.45 ^c
LI	0.22	0.28	0.22	0.20	0.05	-0.19	0.24	0.09	0.25	1.00	0.12	0.11
ткш	-0.82 ^a	-0.22	-0.27	-0.09	-0.45 ^c	-0.28	-0.46 ^c	0.024	0.67 ^b	0.14	1.00	0.10
GY	-0.20	0.36	0.87 ^a	0.76 ^a	0.43 ^c	1.00 ^a	0.41 ^c	0.93 ^a	0.62 ^b	0.20	0.29	1.00

DH=Days to heading (days), DM=Days to maturity (days), PH=Plant height (cm), PL=Panicle length (cm), PdL=Peduncle length (cm), SCD=Second basal culm internode diameter (mm), SPK=No. of spikelet per panicle, TB=Total biomass (t ha 1), GY= Grain yield (t ha 1), LI=Lodging index (%), TKW=Thousand kernel weight (g). a,b, c shows significant association at 0.1,1,5% probability level, respectively.

Table 3: Genotypic (lower diagonal) and phenotypic (upper diagonal) correlation analysis among 12 traits tef breeding lines.

Total biomass showed positive and significant phenotypic and genotypic correlation with almost 60% of the studied traits. Harvest index, on the other hand, showed negative and significant genotypic and phenotypic association withis dailystholytraiding nulet dengthug ty if I and thousaked here and in the observed low [4,MM this study, furthermore, lodging index was not significantly (p=0.05) associated with all studied traits at both phenotypic and genotypic level. Un like this result, a negative and highly significant genotypic association was reported between lodging index and that of culm length, panicle length and plant height in the previous study [5].

Days to heading showed positive and significant genotypic and wh to the set of the second end of the second en total biomass, grain yield, harvest index, lodging index, 1000-kernel weight and grain yield. Thus, it had significant and negative association with harvest index and thousand kernel weight while it showed no significant (P=0.05) association with the remaining five traits. Days to maturity has also exhibited similar association with all traits other than peduncle length, second culm diameter, lodging index, 1000-kernel weight and grain yield at phenotypic level and peduncle length, lodging index, 1000-kernel weight and grain yield at genotypic level (Table 3). Similar results were also reported previously in tef [5,12,13] and in maize [4]. In general, the estimated higher genotypic correlations than their respective phenotypic correlations in this study is in line with the previous reports [4,5]. Besides, the positive and significant association of grain yield with biomass yield and harvest index is also in line with several previous findings [4,5,12,14-17].

Path coefficient analysis

The direct effects of various traits on grain yield ranged from -0.004 to 0.816 at genotypic level (Table 4) and from -0.012 to 0.944 at phenotypic level (Table 5). Thus, total biomass (0.816) followed by harvest index (0.273) and plant height (0.226) showed the highest positive genotypic direct effects on grain yield per plot. Similarly, in the previous studies, the highest direct effect on grain yield were reported for total biomass [3,18] and harvest index [3,5]. Such results suggest to emphasis on the genetic improvement of those traits to increase grain yield by indirect selection [2,3]. In the present study,

number of spikelet per panicle had the highest negative direct effect on grain yield while, the smallest value was exhibited due to days to heading and maturity, peduncle length, second culm diameter and Citation: Habte Jifar Daba (2019) Correlations and Path Coefficient Analysis of Major Quantitative Characters in Tef [*Eragrostis tef* (Zucc.) Trotter] Breeding Lines. Adv Crop Sci Tech 7: 431.

Page 4 of 6

Traits DH DM РН PL Pdl SCD SPK твм нι LI ткw GY -0.085 -0.019 DH -0.079 -0.038 -0.063 0.018 -0.029 -0.078 0.093 -0.019 0.07 -0.2 -0.002 0.36 DM -0.002 -0.002 -0.002 -0.001 -0.002 -0.002 0.001 -0.001 0.001 -0.001 PH 0.1 0.165 0.226 0.231 0.075 0.247 0.183 0.238 -0.032 0.05 -0.061 0.87 PL 0.003 0.003 0.004 0.004 0.005 0.004 0.004 -0.001 0.001 0.00 0.76 0.00 PdL 0.01 -0.011 -0.016 -0.002 -0.048 -0.004 0.006 -0.016 -0.014 -0.002 0.022 0.43 SCD -0.009 -0.011 -0.028 -0.034 -0.002 -0.026 -0.029 -0.037 0.009 0.005 0.007 1.1 SPK -0.13 -0.161 -0.117 0.119 -0.034 0.066 -0.103 -0.116 -0.136 0.019 -0.143 0.41 TBM 0 18 0 547 0 857 0 849 0 278 1 168 0 669 0.816 0 171 0 073 0.016 0.93 -0 297 -0 172 -0.093 0.057 0 273 н -0.038 -0 101 0 0 7 9 -0 226 0.068 0 183 0.62 0.014 0.017 0.014 0.012 0.003 -0.012 0.015 0.006 0.016 0.062 0.009 LI 0.2 TKW 0.018 0.005 0.006 0.002 0.01 0.006 0.01 0.00 -0.014 -0.003 -0.021 0.29

DH=Days to heading (days), DM=Days to maturity (days), PH=Plant height (cm), PL=Panicle length (cm), PdL=Peduncle length (cm), SCD=Second basal culm internode diameter (mm), SPK=No. of spikelet per panicle, TB= Total biomass (t ha⁻¹), GY=Grain yield (t ha⁻¹), LI=Lodging index (%), TKW=Thousand kernel weight (g). Genotypic coefficient of determination = 1.00; Residual variable effect=0.00.

Table 4: Path analysis showing genotypic direct (bold) and indirect effects of various traits on grain yield of tef breeding lines.

Traits	DH	DM	РН	PL	Pdl	SCD	SPK	твм	н	LI	ткw	GY
DH	0.03	0.019	0.011	0.014	-0.004	0.005	0.017	0.006	-0.023	0.004	-0.14	-0.139
DM	-0.018	-0.027	-0.013	-0.012	-0.002	-0.006	-0.011	-0.012	0.011	-0.005	0.21	0.209
PH	0.006	0.007	0.015	0.013	0.004	0.008	0.011	0.014	-0.001	0.003	0.76	0.761
PL	-0.023	-0.022	-0.044	-0.051	-0.001	-0.025	-0.038	-0.038	0.012	-0.007	0.55	0.549
PdL	0.002	-0.001	-0.004	0.00	-0.014	0.00	0.001	-0.004	-0.003	0.00	0.35	0.349
SCD	-0.002	-0.003	-0.006	-0.006	.0.00	-0.012	-0.006	-0.007	0.001	0.001	0.52	0.521
SPK	-0.011	-0.008	-0.014	-0.015	0.001	-0.009	-0.02	-0.013	0.008	-0.003	0.38	0.379
твм	0.189	0.406	0.84	0.698	0.264	0.585	0.594	0.944	0.019	0.028	0.9	0.899
ні	-0.322	-0.17	-0.034	-0.098	0.098	-0.021	-0.178	0.008	0.424	0.059	0.45	0.451
LI	0.006	0.007	0.008	0.005	0.001	-0.004	0.006	0.001	0.006	0.04	0.004	0.119
TKW	0.004	0.001	0.002	0.001	0.002	0.00	0.003	0.00	-0.003	-0.001	-0.011	0.08

DH=Days to heading (days), DM=Days to maturity (days), PH=Plant height (cm), PL=Panicle length (cm), PdL=Peduncle length (cm), SCD=Second basal culm internode diameter (mm), SPK=No. of spikelet per panicle, TB= Total biomass (t ha⁻¹), GY=Grain yield (t ha⁻¹), LI=Lodging index (%), TKW=Thousand kernel weight (g). Phenotypic coefficient of determination = 0.999; Residual variable effect=0.0298.

Table 5: Path analysis showing phenotypic direct (bold) and indirect effects of various traits on grain yield of tef breeding lines.

Principal component analysis

Principal component analysis based on 12 traits of 28 tef breeding lines showed that the first four principal components with eigenvalue greater than one contributed for 81.12% of the total variations. Thus, PC1, 2, 3 and 4 accounted for about 40, 21, 10 and 9.7%, respectively

values of nearly unity at both levels. This indicates that over 99% of the variations in grain yield in the model has been explained through

the independent variables.

(Table 6). In such analysis, plant height followed by panicle length, total biomass and number of spikelet per panicle were the major contributing traits to PC1 while harvest index, days to heading and grain yield were to PC2. Besides, in PC3, the largest contributing traits were 1000-kernel weight, peduncle length and lodging index while that of PC4 were lodging index, peduncle length and second culm

Citation: Habte Jifar Daba (2019) Correlations and Path Coefficient Analysis of Major Quantitative Characters in Tef [*Eragrostis tef* (Zucc.) Trotter] Breeding Lines. Adv Crop Sci Tech 7: 431.

Page 5 of 6

diameter. The total variation obtained in this study is in line with the findings of the previous reports [13,14].

In the PCA-biplot analysis, the studied traits and genotypes were also plotted based on the first two principal components which accounted for over 60% (PC1=40.2% and PC2=20.91%) of the total variations (Figure 1). Based on this biplot, the 28 studied tef genotypes were grouped into four categories with various important traits (Figure 1). Thus, group-I consisted of tef genotypes such as RIL-91, RIL-132, RIL-110 and RIL-271 which had higher values of total biomass, grain yield, peduncle length, lodging index, plant height and second culm diameter. The second group, on the other hand, consisted of genotypes like RIL-171, RIL-85, RIL-69, Quncho, RIL-103, RIL-137, and RIL-96 which had longer days to heading and maturity, large number of spikelet per panicle and longer panicle. The third groups of genotypes include 12 genotypes observed to have high harvest index and 1000-kernel weight. The fourth group of genotypes, however, include RIL-134, RIL-302, RIL-81 and Kinde which had relatively lower values of most of the studied traits. Based on this grouping, genotypes in group-I which have the highest grain yield and total biomass along with lower peduncle length and lodging index have a great potential for the intended yield improvement in tef.

Traits	PC1	PC2	PC3	PC4			
DH	0.24	-0.49	0.01	0.13			
DM	0.27	-0.22	0.11	0.24			
РН	0.43	0.11	-0.01	0.09			
PL	0.40	-0.01	0.13	-0.09			
PdL	0.08	0.25	-0.54	0.46			
SCD	0.29	0.11	-0.03	-0.45			
SPK	0.37	-0.16	0.06	-0.07			
ТВМ	0.41	0.22	-0.04	-0.08			
GY	0.31	0.43	-0.01	0.01			
н	-0.11	0.55	0.06	0.12			
LI	0.07	0.03	0.50	0.66			
ткw	-0.09	0.23	0.65	-0.17			
Eigenvalue	4.82	2.51	1.23	1.17			
Variability (%)	40.20	20.91	10.26	9.74			
Cumulative %	40.20	61.12	71.38	81.12			
DH=Days to heading (days), DM=Days to maturity (days), PH=Plant height (cm), PL=Panicle length (cm), PdL=Peduncle length (cm), SCD=Second basal culm internode diameter (mm), SPK=No. of spikelet per panicle, TB=Total biomass (t ha ⁻¹), GY= Grain yield (t ha ⁻¹), LI=Lodging index (%), TKW=Thousand							

kernel weight (g)

Table 6: Eigenvectors and eigenvalues of the 12 traits of 28 tef breeding lines.

Like the PCA-biplot for genotypes, the PCA-biplot of the 12 studied quantitative traits also showed the formation of three groups of traits whereby group-I, II and III consisted of six, four and two traits, respectively. Thus, group-I consisted of traits such as grain yield, total biomass, plant height, peduncle length, second culm diameter and lodging index which are directly related to yield and lodging. Group-II consisted of days to heading, days to maturity, panicle length and number of spikelet per panicle which are a combination of phenological and morphological traits. The third group, however, consisted of harvest index and 1000-kernel weight.

In such plotting, in general, some groups of traits were found to have acute angle between them while, the other traits have obtuse angle. The nature of this angle is an indication of the degree of association and relationships existing between the different traits. Thus, in the present study, the traits having the smallest angle with each other are known to have strong and positive correlations so as to aid simultaneous improvement of those traits.



Figure 1: PCA-biplot showing the relationship between the studied genotypes (a) and traits (b).

Conclusion

Genotypic correlation analysis in this study showed positive and significant association between grain yield and that of shoot biomass yield per plot, plant height, panicle length, second culm diameter and harvest index. Such kind of association suggests existence of common genetic/physiological basis among those traits to increase their simultaneous improvement. Total biomass and harvest index which correlated positively and significantly with grain yield also exhibited the strongest direct and indirect effects on grain yield at both genotypic and phenotypic level. In general, the results from this study enabled to identify some important traits to be considered in future tef improvement program.

Acknowledgments

The author would like to thank Ethiopian Institute of Agricultural Research for funding the research work. Besides, the technical supports of tef breeding staffs at Holetta and Debre Zeit research center and Alem Tena substation are greatly acknowledged.

References

- 1. CSA (2017) Agricultural sample survey 2016/17 (2009 E.C.) Report on area and production of major crops, (Private Peasant House holdings, Meher Season). Stat Bulletin p: 584.
- 2. Ayalneh T, Amsalu A, Habtamu Z (2012) Genetic divergence, trait association and path analysis of Tef (*Eragrostis tef* (Zucc.) Trotter) Lines. World J Agric Sciences 8: 642-646.

- 3. Nigus C, Mohammed W, Damte T (2016) Genetic variation, correlation and path coefficient analysis in Tef [*Eragrostis tef* (Zucc.) Trotter] Genotypes for yield, yield related traits at Maysiye, Northern Ethiopia. American J Res Communic 4: 73-102.
- Belay N (2018) Genetic variability, heritability, correlation and path coefficient analysis for grain yield and yield component in maize (*Zea* mays L.) Hybrids. Adv Crop Sci Tech 6: 399.
- Lule D, Mengistu G (2014) Correlation and path coefficient analysis of quantitative traits in tef [*Eragrostis tef* (Zucc.) Trotter] germplasm accessions from different regions of Ethiopia. American J Res Commun 2: 194-204.
- Hailu A, Alamerew S, Nigussie M, Assefa E (2016) Correlation and path coefficient analysis of yield and yield associated traits in Barley (*Hordeum vulgare* L.) germplasm. Adv Crop Sci Tech 4: 216.
- Dewey DR, Lu KA (1959) Correlation and path-coefficient analysis of components of crested wheatgrass seed production. Agron J 51: 515-518.
- 8. Gomez KA, Gomez A (1984) Statistical procedures for agricultural research. New York: John Wiley & Sons.
- 9. Alvarado G, Marco L, Mateo V, Angela P, Francisco R, et al. (2016) META-R (Multi Environment Trial Analysis with R for Windows.) Version 6.0, International Maize and Wheat Improvement Center.
- Cruz CD (2016) Genes software-extended and integrated with the R Matlab and Selegen, Acta Scientia rum. Agron 38: 547-552.
- 11. MINITAB (2013) Minitab statistical software, release 17 for windows, State College, Pennsylvania, USA.
- 12. Jifar H, Bekele E, Assefa K (2015) Genetic variability, heritability and association of traits in released tef [*Eragrostis tef* (Zuce.) Trotter] varieties evaluated in Southwestern and Central Ethiopia. J Sci Sust Develop 3: 19-31.
- 13. Jifar H, Tesfaye K, Assefa K, Chanyalew S, Tadele Z (2017) Semi-Dwarf Tef (*Eragrostis tef*) lines for high seed yield and lodging tolerance in Central Ethiopia. African Crop Sci J 25: 419-439.
- 14. Chanyalew S, Tefera H, Zelleke H, Singh H (2006) Correlation and path coefficient analysis of yield related traits in recombinant inbred lines of tef [*Eragrostis tef* (Zucc.) Trotter]. J Genet Breed 60: 209-216.
- Chanyalew S, Tefera H, Singh H (2009) Genetic variability, heritability and trait relationships in recombinant inbred lines of Tef [*Eragrostis tef* (Zucc.) Trotter]. Res J Agric Biol Sci 5: 474-479.
- 16. Lule D, Tesfaye K, Fetene M, De Villiers S (2012) Inheritance and association of quantitative traits in finger millet (Eleusine coracana Subsp. Coracana) landraces collected from Eastern and South Eastern Africa. Int J Genetics 2: 12-22.
- Debebe A, Singh H, Tefera H (2014) Interrelationship and path coefficient analysis of yield components in F4 progenies of Tef [*Eragrostis tef* (Zucc.) Trotter] Pakistan. J Biol Sci 17: 92-97.
- Ayalew H, Genet T, Wondale L (2011) Correlation and path coefficient analysis among yield component traits in Tef [*Eragrostis tef*) Zucc. Trotter] landraces. Libyan Agric Res Cent J Int 2: 180-185.
- 19. Shahid M, Mohammad F, Tahir M (2002) Path coefficient analysis in Wheat. Sarhad J Agric 18: 383-388.